

Flannelmouth sucker Movement: Glen Canyon Dam to Lake Mead, Colorado River

Robin J. Osterhoudt

David L. Rogowski

Arizona Game and Fish Department



Objectives

Flannelmouth sucker:

- Distribution
- Movement
- Growth
- Survival




Photo by Jeff Sorensen

Introduction



- Flannelmouth sucker (*Catostomus latipinnis*) is a native species endemic to the Colorado River basin
- Migration within tributaries occurs during spawning (Paria River, Little Colorado River, Havasu creek, others)
- Live >15 years, can exceed 600 mm total length
- Highly fecund (15,000 eggs/year) and may spawn for more than 2 decades


Study Area



Methods







- Electrofishing, hoop nets, trammel nets, seining throughout Colorado River by various agencies (AGFD, USFWS, GCMRC, NPS)
- PIT tag flannelmouth sucker (FMS) ≥ 150 mm total length (TL)
- Filtered data for FMS ≥ 150 mm TL that have been recaptured at least once (4,087 records)
- Randomized PIT tag numbers, sub-sampled 212 unique FMS
- Measures:
 - Average distance moved (miles) between encounters over time (1 week)
 - Growth (mm)/year for 168 fish (adjusted for errors)
 - Apparent yearly survival between years using program Mark

FMS Distribution (Preliminary data)



FMS Catch-per-unit-effort (CPUE; fish/hr.) by Reach 2000-2014 (Preliminary data)


CPUE (fish/hr)



Year

Results (Preliminary data)

Individual FMS Movement per Week N= 212



Results (Preliminary data)

43% of FMS visited the Little Colorado River!



Results


(Preliminary data)

Size class (mm; TL)	N	Average growth mm/year	95% CI
All FMS	168	47	[41.84, 52.23]
150-350	111	59.7	[53.41, 66.00]
>350	57	22.4	[17.64, 27.09]



Robin J. Osterhoudt

FMS Apparent Yearly Survival (Preliminary data)



Conclusions

Flannelmouth sucker

•Movement:

- Have high site fidelity, moving very little between recaptures (~10 miles/week)
- Can move great distances (140+ miles/week)
- Frequently found within the LCR (43%)

•Growth:

- Overall average 47 mm/year
- Younger fish (150-350 mm TL) grow more than fish >350 mm TL (59 mm/year vs. 22 mm/year)

•Survival:

- 42% over 12 years with some variation

Future Work

- Evaluate more tagged fish
- How does FMS movement relate to environmental factors (flow and water temperature)?



Robin J. Osterhoudt




Robin J. Osterhoudt

Acknowledgements



FMS Length Frequency of Captured Fish (Preliminary data)



FMS Yearly Capture Probability (Preliminary data)

